

The Kastler-Kalau-Walze type theorem for 6-dimensional manifolds with boundary

Jian Wang, Yong Wang*

School of Mathematics and Statistics, Northeast Normal University, Changchun, 130024, P.R.China

Abstract

In this paper, we define lower dimensional volumes of spin manifolds with boundary. We compute the lower dimensional volume $\text{Vol}_6^{(1,3)}$ for 6-dimensional spin manifolds with boundary and the gravity on boundary is derived by the noncommutative residue associated with Dirac operators. For 6-dimensional manifolds with boundary, we also get a Kastler-Kalau-Walze type theorem for a general fourth order operator .

Keywords: lower-dimensional volumes; noncommutative residue; gravitational action; perturbations of Dirac operators.

2000 MSC: 53G20, 53A30, 46L87

1. Introduction

The noncommutative residue plays a prominent role in noncommutative geometry [1][2]. Connes [3] used the noncommutative residue to derive a conformal 4-dimensional Polyakov action analogy. Connes [4] proved that the noncommutative residue on a compact manifold M coincided with the Dixmier's trace on pseudodifferential operators of order $-\dim M$. Several years ago, Connes made a challenging observation that the noncommutative residue of the square of the inverse of the Dirac operator was proportional to the Einstein-Hilbert action, which we call the Kastler-Kalau-Walze theorem. Kastler[5] gave a brute-force proof of this theorem. Kalau and Walze [6] proved this theorem in the normal coordinates system simultaneously. Ackermann [7] gave a note on a new proof of this theorem by means of the heat kernel expansion.

Recently, Ponge defined lower dimensional volumes of Riemannian manifolds by the Wodzicki residue [8]. Fedosov et al defined a noncommutative residue on Boutet de Monvel's algebra and proved that it was a unique continuous trace [9]. Wang generalized the Connes' results to the case of manifolds with boundary in [10] [11], and proved a Kastler-Kalau-Walze type theorem for the Dirac operator and the signature operator for lower-dimensional manifolds with boundary[12] [13]. The purpose of papers [12] [13] is to derive the gravitational action by the noncommutative residue associated with Dirac operators for spin manifolds with boundary, but the boundary term vanished in Wang's results. The motivation of this paper is to derive the gravitational action on boundary by the noncommutative residue associated with Dirac operators. In other words, we want to get a nonvanishing boundary term. In [12] [13], Wang computed $\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{-1}]$ and $\widetilde{\text{Wres}}[\pi^+ D^{-2} \circ \pi^+ D^{-2}]$, where the two operators are symmetric. In this paper, for 6-dimensional manifolds with boundary, we compute $\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{-3}]$. Since D^{-1} and D^{-3} are not symmetric, the boundary term is the integral of the extrinsic scalar curvature and the gravitational action on boundary emerges. We also get a generalized Kastler-Kalau-Walze theorem associated with a general fourth order operators for 6-dimensional spin manifolds.

This paper is organized as follows: In Section 2, we define lower dimensional volumes of spin manifolds with boundary. In Section 3, for 6-dimensional spin manifolds with boundary and the associated Dirac

*Corresponding author; Email address: wangy581@nenu.edu.cn (Yong Wang)

Email address: wangj068@gmail.com (Jian Wang)

operator D and D^3 , we compute the lower dimensional volume $\text{Vol}_6^{(1,3)}$ and get a Kastler-Kalau-Walze type theorem in this case. In Section 4, we get a Kastler-Kalau-Walze type theorem associated with the fourth order operators for 6-dimensional spin manifolds.

2. Lower dimensional volumes of spin manifolds with boundary

In order to define lower dimensional volumes of spin manifolds with boundary, we need some basic facts and formulae about Boutet de Monvel's calculus and the definition of the noncommutative residue for manifolds with boundary. We can find them in Section 2,3 [11] and Section 2.1[12].

Let M be a n -dimensional compact oriented spin manifold with boundary ∂M . We assume that the metric g^M on M has the following form near the boundary,

$$g^M = \frac{1}{h(x_n)} g^{\partial M} + dx_n^2, \quad (2.1)$$

where $g^{\partial M}$ is the metric on ∂M . Let D be the Dirac operator associated to g on the spinors bundle $S(TM)$ [12]. Let p_1, p_2 be nonnegative integers and $p_1 + p_2 \leq n$.

Definition 2.1. *Lower dimensional volumes of spin manifolds with boundary are defined by*

$$\text{Vol}_n^{(p_1, p_2)} M := \widetilde{\text{Wres}}[\pi^+ D^{-p_1} \circ \pi^+ D^{-p_2}] \quad (2.2)$$

where the related definitions, see Section 2, 3[12].

Denote by $\sigma_l(A)$ the l -order symbol of an operator A . By (2.1.4)-(2.1.8)[12], we get

$$\widetilde{\text{Wres}}[\pi^+ D^{-p_1} \circ \pi^+ D^{-p_2}] = \int_M \int_{|\xi|=1} \text{trace}_{S(TM)}[\sigma_{-n}(D^{-p_1-p_2})] \sigma(\xi) dx + \int_{\partial M} \Phi, \quad (2.3)$$

and

$$\begin{aligned} \Phi &= \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{j,k=0}^{\infty} \sum \frac{(-i)^{|\alpha|+j+k+1}}{\alpha!(j+k+1)!} \times \text{trace}_{S(TM)}[\partial_{x_n}^j \partial_{\xi'}^\alpha \partial_{\xi_n}^k \sigma_r^+(D^{-p_1})(x', 0, \xi', \xi_n) \\ &\quad \times \partial_{x'}^\alpha \partial_{\xi_n}^{j+1} \partial_{x_n}^k \sigma_l(D^{-p_2})(x', 0, \xi', \xi_n)] d\xi_n \sigma(\xi') dx', \end{aligned} \quad (2.4)$$

where the sum is taken over $r - k - |\alpha| + l - j - 1 = -n$, $r \leq -p_1, l \leq -p_2$.

Since $[\sigma_{-n}(D^{-p_1-p_2})]|_M$ has the same expression as $\sigma_{-n}(D^{-p_1-p_2})$ in the case of manifolds without boundary, so locally we can use the computations [5], [6], [8], [12] to compute the first term. The following proposition is the motivation of the definition of lower dimensional volumes of spin manifolds with boundary [13].

Proposition 2.2. [13] *Lower dimensional volumes of spin manifolds with boundary are given by*

- 1) When $p_1 + p_2 = n$, then $\text{Vol}_n^{(p_1, p_2)} M = c_0 \text{Vol}_M$.
- 2) when $p_1 + p_2 \equiv n \pmod{1}$, $\text{Vol}_n^{(p_1, p_2)} M = \int_{\partial M} \Phi$.
- 3)

$$\text{Vol}_4^{(1,1)} = -\frac{\Omega_4}{3} \int_M s \text{dvol}_M; \quad \text{Vol}_3^{(1,1)} = c_1 \text{Vol}_{\partial M} \quad (2.5)$$

where c_0, c_1 are constants and s is the scalar curvature.

3. A Kastler-Kalau-Walze type theorem for 6-dimensional spin manifolds with boundary

In this section, We compute the lower dimensional volume $\text{Vol}_6^{(1,3)}$ for 6-dimensional spin manifolds with boundary and get a Kastler-Kalau-Walze type theorem in this case.

Firstly, we compute the symbol $\sigma(D^{-3})$ of D^{-3} . Recall the definition of the Dirac operator D [9][14]. Let ∇^L denote the Levi-Civita connection about g^M . In the local coordinates $\{x_i; 1 \leq i \leq n\}$ and the fixed orthonormal frame $\{\tilde{e}_1, \dots, \tilde{e}_n\}$, the connection matrix $(\omega_{s,t})$ is defined by

$$\nabla^L(\tilde{e}_1, \dots, \tilde{e}_n) = (\tilde{e}_1, \dots, \tilde{e}_n)(\omega_{s,t}). \quad (3.1)$$

The Dirac operator is defined by

$$D = \sum_{i=1}^n c(\tilde{e}_i) \left[\tilde{e}_i - \frac{1}{4} \sum_{s,t} \omega_{s,t}(\tilde{e}_i) c(\tilde{e}_s) c(\tilde{e}_t) \right]. \quad (3.2)$$

where $c(\tilde{e}_i)$ denotes the Clifford action.

Recall the definition of the Dirac operator D^2 in [5], [6] and [13], we have

$$D^2 = - \sum_{i,j} g^{i,j} \left[\partial_i \partial_j + 2\sigma_i \partial_j + (\partial_i \sigma_j) + \sigma_i \sigma_j - \Gamma_{i,j}^k \partial_i - \Gamma_{i,j}^k \sigma_k \right] + \frac{1}{4} s. \quad (3.3)$$

where $\sigma_i := -\frac{1}{4} \sum_{s,t} \omega_{s,t}(\partial_i) e_s e_t$.

Combining (3.2) and (3.3), we have

$$\begin{aligned} D^3 &= \sum_{i=1}^n c(\tilde{e}_i) \langle e_i, dx_i \rangle \left\{ \left(- \sum_{i,j} g^{i,j} \partial_i \partial_j \right) \partial_l - \sum_{i,j} \partial_l (g^{i,j}) \partial_i \partial_j \right. \\ &\quad + \left[- \sum_{i,j} g^{i,j} (2\sigma_i \partial_j - \Gamma_{i,j}^k \partial_i) \right] \partial_l - \sum_{i,j} \partial_l (g^{i,j}) (2\sigma_i \partial_j - \Gamma_{i,j}^k \partial_i) \\ &\quad - \sum_{i,j} g^{i,j} \left(2(\partial_l \sigma_i) \partial_j - \partial_l (\Gamma_{i,j}^k) \partial_i \right) \\ &\quad + \left[- \sum_{i,j} g^{i,j} \left((\partial_i \sigma_j) + \sigma_i \sigma_j - \Gamma_{i,j}^k \sigma_k \right) + \frac{1}{4} s \right] \partial_l \\ &\quad - \sum_{i,j} \partial_l (g^{i,j}) \left((\partial_i \sigma_j) + \sigma_i \sigma_j - \Gamma_{i,j}^k \sigma_k \right) + \frac{1}{4} \partial_l s \\ &\quad \left. - \sum_{i,j} g^{i,j} \partial_l \left((\partial_i \sigma_j) + \sigma_i \sigma_j - \Gamma_{i,j}^k \sigma_k \right) + \frac{1}{4} \partial_l s \right\} \\ &\quad - \frac{1}{4} \sum_{s,t} \omega_{s,t}(\tilde{e}_l) c(e_l) c(\tilde{e}_s) c(\tilde{e}_t) \left\{ - \sum_{i,j} g^{i,j} \partial_i \partial_j - \sum_{i,j} g^{i,j} (2\sigma_i \partial_j - \Gamma_{i,j}^k \partial_i) \right. \\ &\quad \left. - \sum_{i,j} g^{i,j} \left((\partial_i \sigma_j) + \sigma_i \sigma_j - \Gamma_{i,j}^k \sigma_k \right) + \frac{1}{4} s \right\}. \end{aligned} \quad (3.4)$$

Then, we obtain

$$\sigma_3(D^3) = \sqrt{-1}c(\xi)|\xi|^2, \quad (3.5)$$

$$\sigma_2(D^3) = c(\xi)(2\sigma^k - \Gamma^k)\xi_k - \frac{1}{4}|\xi|^2 \sum_{s,t} \omega_{s,t}(\tilde{e}_l)c(e_l)c(\tilde{e}_s)c(\tilde{e}_t), \quad (3.6)$$

$$\begin{aligned} \sigma_1(D^3) &= \sum_{i=1}^n c(\tilde{e}_l)\langle e_l, dx_l \rangle \left\{ - \sum_{i,j} \partial_l(g^{i,j})(2\sigma_i\partial_j - \Gamma_{i,j}^k\partial_i) \right. \\ &\quad + \sum_{i,j} g^{i,j}\partial_l(\Gamma_{i,j}^k)\partial_i - \sum_{i,j} g^{i,j}\left(2(\partial_l\sigma_i)\partial_j\right) \\ &\quad + \left[- \sum_{i,j} g^{i,j}\left((\partial_i\sigma_j) + \sigma_i\sigma_j - \Gamma_{i,j}^k\sigma_k\right) + \frac{1}{4}s \right] \partial_l \Big\} \\ &\quad - \frac{1}{4} \sum_{s,t} \omega_{s,t}(\tilde{e}_l)c(e_l)c(\tilde{e}_s)c(\tilde{e}_t) \left[- \sum_{i,j} g^{i,j}(2\sigma_i\partial_j - \Gamma_{i,j}^k\partial_i) \right], \end{aligned} \quad (3.7)$$

$$\begin{aligned} \sigma_0(D^3) &= \sum_{i=1}^n c(\tilde{e}_l)\langle e_l, dx_l \rangle \left\{ - \sum_{i,j} \partial_l(g^{i,j})\left((\partial_i\sigma_j) + \sigma_i\sigma_j - \Gamma_{i,j}^k\sigma_k\right) + \frac{1}{4}\partial_l s \right. \\ &\quad - \sum_{i,j} g^{i,j}\partial_l\left((\partial_i\sigma_j) + \sigma_i\sigma_j - \Gamma_{i,j}^k\sigma_k\right) + \frac{1}{4}\partial_l s \Big\} \\ &\quad - \frac{1}{4} \sum_{s,t} \omega_{s,t}(\tilde{e}_l)c(e_l)c(\tilde{e}_s)c(\tilde{e}_t) \left[- \sum_{i,j} g^{i,j}\left((\partial_i\sigma_j) + \sigma_i\sigma_j - \Gamma_{i,j}^k\sigma_k\right) + \frac{1}{4}s \right]. \end{aligned} \quad (3.8)$$

Write

$$D_x^\alpha = (-\sqrt{-1})^{|\alpha|}\partial_x^\alpha; \quad \sigma(D^3) = p_3 + p_2 + p_1 + p_0; \quad \sigma(D^{-3}) = \sum_{j=3}^{\infty} q_{-j}. \quad (3.9)$$

By the composition formula of psudodifferential operators, we have

$$\begin{aligned} 1 = \sigma(D^3 \circ D^{-3}) &= \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} [\sigma(D)] D_x^{\alpha} [\sigma(D^{-1})] \\ &= (p_3 + p_2 + p_1 + p_0)(q_{-3} + q_{-4} + q_{-5} + \cdots) \\ &\quad + \sum_j (\partial_{\xi_j} p_3 + \partial_{\xi_j} p_2 + \partial_{\xi_j} p_1 + \partial_{\xi_j} p_0)(D_{x_j} q_{-3} + D_{x_j} q_{-4} + D_{x_j} q_{-5} + \cdots) \\ &= p_3 q_{-3} + (p_3 q_{-4} + p_2 q_{-3} + \sum_j \partial_{\xi_j} p_3 D_{x_j} q_{-3}) + \cdots, \end{aligned} \quad (3.10)$$

Then we obtain

$$q_{-3} = p_3^{-1}; \quad q_{-4} = -p_3^{-1}[p_2 p_3^{-1} + \sum_j \partial_{\xi_j} p_3 D_{x_j}(p_3^{-1})]. \quad (3.11)$$

By Lemma 2.1 in [12] and (3.4)-(3.11), we obtain

Lemma 3.1.

$$\sigma_{-1}(D^{-1}) = \frac{\sqrt{-1}c(\xi)}{|\xi|^2}; \quad (3.12)$$

$$\sigma_{-2}(D^{-1}) = \frac{c(\xi)\sigma_0(D)c(\xi)}{|\xi|^4} + \frac{c(\xi)}{|\xi|^6} \sum_j c(dx_j) \left(\partial_{x_j}[c(\xi)]|\xi|^2 - c(\xi)\partial_{x_j}(|\xi|^2) \right); \quad (3.13)$$

$$\sigma_{-3}(D^{-3}) = \frac{\sqrt{-1}c(\xi)}{|\xi|^4}; \quad (3.14)$$

$$\begin{aligned} \sigma_{-4}(D^{-3}) &= \frac{c(\xi)\sigma_2(D^3)c(\xi)}{|\xi|^8} + \frac{\sqrt{-1}c(\xi)}{|\xi|^8} \left(|\xi|^4 c(dx_n)\partial_{x_n}c(\xi') - 2h'(0)c(dx_n)c(\xi) \right. \\ &\quad \left. + 2\xi_n c(\xi)\partial_{x_n}c(\xi') + 4\xi_n h'(0) \right), \end{aligned} \quad (3.15)$$

where $\sigma_0(D) = -\frac{1}{4} \sum_{s,t} \omega_{s,t}(\tilde{e}_i)c(\tilde{e}_i)c(\tilde{e}_s)c(\tilde{e}_t)$.

Since Φ is a global form on ∂M , so for any fixed point $x_0 \in \partial M$, we can choose the normal coordinates U of x_0 in ∂M (not in M) and compute $\Phi(x_0)$ in the coordinates $\tilde{U} = U \times [0, 1) \subset M$ and the metric $\frac{1}{h(x_n)}g^{\partial M} + dx_n^2$. The dual metric of g^M on $\tilde{U}$ is $h(x_n)g^{\partial M} + dx_n^2$. Write $g_{ij}^M = g^M(\frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j})$; $g_M^{ij} = g^M(dx_i, dx_j)$, then

$$[g_{i,j}^M] = \begin{bmatrix} \frac{1}{h(x_n)}[g_{i,j}^{\partial M}] & 0 \\ 0 & 1 \end{bmatrix}; \quad [g_M^{i,j}] = \begin{bmatrix} h(x_n)[g_{\partial M}^{i,j}] & 0 \\ 0 & 1 \end{bmatrix}, \quad (3.16)$$

and

$$\partial_{x_s} g_{ij}^{\partial M}(x_0) = 0, 1 \leq i, j \leq n-1; \quad g_{ij}^M(x_0) = \delta_{ij}. \quad (3.17)$$

Let $n = 6$ and $\{e_1, \dots, e_{n-1}\}$ be an orthonormal frame field in U about $g^{\partial M}$ which is parallel along geodesics and $e_i(x_0) = \frac{\partial}{\partial x_i}(x_0)$, then $\{\tilde{e}_1 = \sqrt{h(x_n)}e_1, \dots, \tilde{e}_{n-1} = \sqrt{h(x_n)}e_{n-1}, \tilde{e}_n = dx_n\}$ is the orthonormal frame field in $\tilde{U}$ about g^M . Locally $S(TM)|_{\tilde{U}} \cong \tilde{U} \times \wedge_{\mathbf{C}}^*(\frac{n}{2})$. Let $\{f_1, \dots, f_8\}$ be the orthonormal basis of $\wedge_{\mathbf{C}}^*(\frac{n}{2})$. Take a spin frame field $\sigma : \tilde{U} \rightarrow \text{Spin}(M)$ such that $\pi\sigma = \{\tilde{e}_1, \dots, \tilde{e}_n\}$, where $\pi : \text{Spin}(M) \rightarrow O(M)$ is a double covering, then $\{[(\sigma, f_i)], 1 \leq i \leq 8\}$ is an orthonormal frame of $S(TM)|_{\tilde{U}}$. In the following, since the global form Φ is independent of the choice of the local frame, so we can compute $\text{tr}_{S(TM)}$ in the frame $\{[(\sigma, f_i)], 1 \leq i \leq 8\}$. Let $\{E_1, \dots, E_n\}$ be the canonical basis of $\mathbf{R}^n$ and $c(E_i) \in \text{cl}_{\mathbf{C}}(n) \cong \text{Hom}(\wedge_{\mathbf{C}}^*(\frac{n}{2}), \wedge_{\mathbf{C}}^*(\frac{n}{2}))$ be the Clifford action. By [Y], then

$$c(\tilde{e}_i) = [(\sigma, c(E_i))]; \quad c(\tilde{e}_i)[(\sigma, f_i)] = [(\sigma, c(E_i)f_i)]; \quad \frac{\partial}{\partial x_i} = [(\sigma, \frac{\partial}{\partial x_i})], \quad (3.18)$$

then we have $\frac{\partial}{\partial x_i}c(\tilde{e}_i) = 0$ in the above frame. By Lemma 2.2 in [12], we have

Lemma 3.2. [12] *For n -dimensional spin manifolds with boundary,*

$$\partial_{x_j}(|\xi|_{g^M}^2)(x_0) = 0, \text{ if } j < n; \quad = h'(0)|\xi'|_{g^{\partial M}}^2, \text{ if } j = n. \quad (3.19)$$

$$\partial_{x_j}[c(\xi)](x_0) = 0, \text{ if } j < n; \quad = \partial_{x_n}[c(\xi')](x_0), \text{ if } j = n, \quad (3.20)$$

where $\xi = \xi' + \xi_n dx_n$.

Next we compute $\sigma_0(D)(x_0)$. By Lemma 2.3 in [12], we have

Lemma 3.3. [12] *For 6-dimensional spin manifolds with boundary,*

$$\sigma_0(D)(x_0) = -\frac{5}{4}h'(0)c(dx_n). \quad (3.21)$$

Now we can compute Φ (see formula (2.4) for the definition of Φ), since the sum is taken over $-r-l+k+j+|\alpha|=5$, $r \leq -1, l \leq -4$, then we have the following five cases:

case a) I) $r = -1, l = -3, k = j = 0, |\alpha| = 1$

From (2.4) we have

$$\text{case a) I)} = - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{|\alpha|=1} \text{trace}[\partial_{\xi'}^{\alpha} \pi_{\xi_n}^{+} \sigma_{-1}(D^{-1}) \times \partial_{x'}^{\alpha} \partial_{\xi_n} \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \quad (3.22)$$

By Lemma 3.2, for $i < n$, then

$$\begin{aligned} \partial_{x_i} \sigma_{-3}(D^{-3})(x_0) &= \partial_{x_i} \left[\frac{\sqrt{-1} c(\xi)}{|\xi|^4} \right](x_0) \\ &= \frac{\sqrt{-1} \partial_{x_i} [c(\xi)](x_0)}{|\xi|^4} - \frac{2\sqrt{-1} c(\xi) \partial_{x_i} [|\xi|^2](x_0)}{|\xi|^6} = 0. \end{aligned} \quad (3.23)$$

Then case a) I) vanishes.

case a) II) $r = -1, l = -3, k = |\alpha| = 0, j = 1$

From (2.4) we have

$$\text{case a) II)} = -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{x_n} \pi_{\xi_n}^{+} \sigma_{-1}(D^{-1}) \times \partial_{\xi_n}^2 \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \quad (3.24)$$

By (2.2.23) in [12], we have

$$\pi_{\xi_n}^{+} \partial_{x_n} \sigma_{-1}(D^{-1})(x_0)|_{|\xi'|=1} = \frac{\partial_{x_n} [c(\xi')](x_0)}{2(\xi_n - i)} + \sqrt{-1} h'(0) \left[\frac{ic(\xi')}{4(\xi_n - i)} + \frac{c(\xi') + ic(dx_n)}{4(\xi_n - i)^2} \right]. \quad (3.25)$$

By (3.14), we obtain

$$\partial_{\xi_n}^2 \sigma_{-3}(D^{-3}) = \sqrt{-1} \left(\frac{(20\xi_n^2 - 4)c(\xi') + 12(\xi_n^3 - \xi_n)c(dx_n)}{(1 + \xi_n^2)^4} \right); \quad (3.26)$$

Since $n = 6$, $\text{tr}_{S(TM)}[\text{id}] = \dim(\wedge^*(3)) = 8$. By the relation of the Clifford action and $\text{tr}AB = \text{tr}BA$, then we have the equalities:

$$\begin{aligned} \text{tr}[c(\xi')c(dx_n)] &= 0; \quad \text{tr}[c(dx_n)^2] = -8; \quad \text{tr}[c(\xi')^2](x_0)|_{|\xi'|=1} = -8; \\ \text{tr}[\partial_{x_n} c(\xi')c(dx_n)] &= 0; \quad \text{tr}[\partial_{x_n} c(\xi')c(\xi')](x_0)|_{|\xi'|=1} = -4h'(0). \end{aligned} \quad (3.27)$$

By (3.25)-(3.27) and direct computations, we have

$$\begin{aligned} &\text{tr} \left\{ \left[\frac{\partial_{x_n} [c(\xi')](x_0)}{2(\xi_n - i)} + \sqrt{-1} h'(0) \left(\frac{ic(\xi')}{4(\xi_n - i)} + \frac{c(\xi') + ic(dx_n)}{4(\xi_n - i)^2} \right) \right] \right. \\ &\quad \times \left. \left[\sqrt{-1} \left(\frac{(20\xi_n^2 - 4)c(\xi') + 12(\xi_n^3 - \xi_n)c(dx_n)}{(1 + \xi_n^2)^4} \right) \right] \right\} (x_0)|_{|\xi'|=1} \\ &= h'(0) \frac{-8 - 24i\xi_n + 40\xi_n^2 + 24i\xi_n^3}{(\xi_n - i)^6 (\xi_n + i)^4}. \end{aligned} \quad (3.28)$$

By (3.24) and (3.28), we obtain

$$\begin{aligned} \text{case a) II)} &= \int_{|\xi'|=1} \int_{-\infty}^{+\infty} h'(0) \frac{4 + 12i\xi_n - 20\xi_n^2 - 12i\xi_n^3}{(\xi_n - i)^6 (\xi_n + i)^4} d\xi_n \sigma(\xi') dx' \\ &= h'(0) \Omega_4 \int_{\Gamma^+} \frac{4 + 12i\xi_n - 20\xi_n^2 - 12i\xi_n^3}{(\xi_n - i)^6 (\xi_n + i)^4} d\xi_n dx' \\ &= h'(0) \Omega_4 \frac{2\pi i}{5!} \left[\frac{4 + 12i\xi_n - 20\xi_n^2 - 12i\xi_n^3}{(\xi_n + i)^4} \right]^{(5)} \Big|_{\xi_n=i} dx' \\ &= -\frac{15}{16} \pi h'(0) \Omega_4 dx', \end{aligned} \quad (3.29)$$

where Ω_4 is the canonical volume of S^4 .

case a) III) $r = -1$, $l = -3$ $j = |\alpha| = 0$, $k = 1$

From (2.4) we have

$$\text{case a) III)} = -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \quad (3.30)$$

By (2.2.29) in [12], we have

$$\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D^{-1})(x_0)|_{|\xi'|=1} = -\frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)^2}. \quad (3.31)$$

By (3.14), we obtain

$$\partial_{\xi_n} \partial_{x_n} \sigma_{-3}(D^{-3})(x_0)|_{|\xi'|=1} = -2\sqrt{-1}h'(0) \left[\frac{(1 - 5\xi_n^2)c(dx_n)}{(1 + \xi_n^2)^4} - \frac{6\xi_n c(\xi')}{(1 + \xi_n^2)^4} \right] - \frac{4\sqrt{-1}\xi_n \partial_{x_n} c(\xi')(x_0)}{(1 + \xi_n^2)^3}. \quad (3.32)$$

By (3.27), (3.31) and (3.32), we obtain

$$\begin{aligned} & \text{tr} \left\{ \left[\frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)^2} \right] \times \left[2\sqrt{-1}h'(0) \left(\frac{(1 - 5\xi_n^2)c(dx_n)}{(1 + \xi_n^2)^4} - \frac{6\xi_n c(\xi')}{(1 + \xi_n^2)^4} \right) + \frac{4\sqrt{-1}\xi_n \partial_{x_n} c(\xi')(x_0)}{(1 + \xi_n^2)^3} \right] \right\} (x_0)|_{|\xi'|=1} \\ &= h'(0) \frac{8i - 32\xi_n - 8i\xi_n^2}{(\xi_n - i)^5(\xi_n + i)^4}. \end{aligned} \quad (3.33)$$

Then

$$\begin{aligned} \text{case a) III)} &= h'(0) \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{-4i + 16\xi_n + 4i\xi_n^2}{(\xi_n - i)^5(\xi_n + i)^4} d\xi_n \sigma(\xi') dx' \\ &= h'(0) \frac{2\pi i}{4!} \left[\frac{-4i + 16\xi_n + 4i\xi_n^2}{(\xi_n + i)^4} \right]^{(4)}|_{\xi_n=i} \Omega_4 dx' \\ &= \frac{25}{16} \pi h'(0) \Omega_4 dx'. \end{aligned} \quad (3.34)$$

case b) $r = -1$, $l = -4$, $k = j = |\alpha| = 0$

From (2.4) and an integration by parts, we have

$$\begin{aligned} \text{case b)} &= -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \times \partial_{\xi_n} \sigma_{-4}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx' \\ &= i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \times \sigma_{-4}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \end{aligned} \quad (3.35)$$

In the normal coordinate, $g^{ij}(x_0) = \delta_i^j$ and $\partial_{x_j}(g^{\alpha\beta})(x_0) = 0$, if $j < n$; $= h'(0)\delta_\beta^\alpha$, if $j = n$. So by Lemma A.2 in [12], we have $\Gamma^n(x_0) = \frac{5}{2}h'(0)$ and $\Gamma^k(x_0) = 0$ for $k < n$. By the definition of δ^k and Lemma 2.3 in

[12], we have $\delta^n(x_0) = 0$ and $\delta^k = \frac{1}{4}h'(0)c(\tilde{e}_k)c(\tilde{e}_n)$ for $k < n$. Then by (3.15), we obtain

$$\begin{aligned}
\sigma_{-4}(D^{-3}) &= \frac{c(\xi)\sigma_2(D^3)c(\xi)}{|\xi|^8} - \frac{c(\xi)}{|\xi|^4} \sum_j \partial_{\xi_j} [c(\xi)|\xi|^2] D_{x_j} \left[\frac{\sqrt{-1}c(\xi)}{|\xi|^4} \right] \\
&= \frac{1}{|\xi|^8} c(\xi) \left(\frac{1}{2} h'(0) c(\xi) \sum_{k < n} \xi_k c(\tilde{e}_k) c(\tilde{e}_n) - \frac{5}{2} h'(0) \xi_n c(\xi) - \frac{5}{4} h'(0) |\xi|^2 c(dx_n) \right) c(\xi) \\
&\quad + \frac{\sqrt{-1}c(\xi)}{|\xi|^8} \left(|\xi|^4 c(dx_n) \partial_{x_n} c(\xi') - 2h'(0) c(dx_n) c(\xi) + 2\xi_n c(\xi) \partial_{x_n} c(\xi') + 4\xi_n h'(0) \right) \\
&= \frac{1}{|\xi|^8} c(\xi) \left(\frac{1}{2} h'(0) c(\xi) c(\xi') c(dx_n) - \frac{5}{2} h'(0) \xi_n c(\xi) - \frac{5}{4} h'(0) |\xi|^2 c(dx_n) \right) c(\xi) \\
&\quad + \frac{1}{|\xi|^8} \left(i|\xi|^4 c(\xi) c(dx_n) \partial_{x_n} c(\xi') - 2ih'(0) c(\xi) c(dx_n) c(\xi) \right. \\
&\quad \left. + 2i\xi_n c(\xi) c(\xi) \partial_{x_n} c(\xi') + 4i\xi_n h'(0) c(\xi) \right) \\
&= \frac{1}{(1 + \xi_n^2)^4} \left(\left(\frac{11}{2} \xi_n (1 + \xi_n^2) + 8i\xi_n \right) h'(0) c(\xi') \right. \\
&\quad \left. + \left(-2i + 6i\xi_n^2 - \frac{7}{4}(1 + \xi_n^2) + \frac{15}{4}\xi_n^2(1 + \xi_n^2) \right) h'(0) c(dx_n) \right. \\
&\quad \left. - 3i\xi_n(1 + \xi_n^2) \partial_{x_n} c(\xi') + i(1 + \xi_n^2) c(\xi') c(dx_n) \partial_{x_n} c(\xi') \right). \tag{3.36}
\end{aligned}$$

By (3.27), (3.31) and (3.36), we obtain

$$\begin{aligned}
&\text{tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \times \sigma_{-4}(D^{-3})] \\
&= \text{tr} \left[-\frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)^2} \times \frac{1}{(1 + \xi_n^2)^4} \left(\left(\frac{11}{2} \xi_n (1 + \xi_n^2) + 8i\xi_n \right) h'(0) c(\xi') \right. \right. \\
&\quad \left. \left. + \left(-2i + 6i\xi_n^2 - \frac{7}{4}(1 + \xi_n^2) + \frac{15}{4}\xi_n^2(1 + \xi_n^2) \right) h'(0) c(dx_n) \right. \right. \\
&\quad \left. \left. - 3i\xi_n(1 + \xi_n^2) \partial_{x_n} c(\xi') + i(1 + \xi_n^2) c(\xi') c(dx_n) \partial_{x_n} c(\xi') \right) \right] \\
&= h'(0) \frac{7 + 6i - (20 - 15i)\xi_n + (7 - 6i)\xi_n^2 + 15i\xi_n^3}{(\xi_n - i)^5(\xi_n + i)^4}. \tag{3.37}
\end{aligned}$$

Then

$$\begin{aligned}
\text{case b)} &= ih'(0) \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{7 + 6i - (20 - 15i)\xi_n + (7 - 6i)\xi_n^2 + 15i\xi_n^3}{(\xi_n - i)^5(\xi_n + i)^4} d\xi_n \sigma(\xi') dx' \\
&= ih'(0) \frac{2\pi i}{4!} \left[\frac{7 + 6i - (20 - 15i)\xi_n + (7 - 6i)\xi_n^2 + 15i\xi_n^3}{(\xi_n + i)^4} \right]^{(4)} \Big|_{\xi_n=i} \Omega_4 dx' \\
&= \left(-\frac{25}{8} - \frac{35i}{16} \right) \pi h'(0) \Omega_4 dx'. \tag{3.38}
\end{aligned}$$

case c) $r = -2$, $l = -3$, $k = j = |\alpha| = 0$

From (2.4) we have

$$\text{case c)} = -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\pi_{\xi_n}^+ \sigma_{-2}(D^{-1}) \times \partial_{\xi_n} \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \tag{3.39}$$

By (2.2.34) , (2.2.37) and (2.2.40) in [13], we have

$$\begin{aligned}
\pi_{\xi_n}^+ \sigma_{-2}(D^{-1})(x_0)|_{|\xi'|=1} &= \pi_{\xi_n}^+ \left[\frac{c(\xi)p_0(x_0)c(\xi) + c(\xi)c(dx_n)\partial_{x_n}[c(\xi')](x_0)}{(1+\xi_n^2)^2} \right] \\
&\quad - h'(0)\pi_{\xi_n}^+ \left[\frac{c(\xi)c(dx_n)c(\xi)}{(1+\xi_n)^3} \right] \\
&:= B_1 - B_2.
\end{aligned} \tag{3.40}$$

where

$$\begin{aligned}
B_1 &= \frac{-1}{4(\xi_n - i)^2} [(2 + i\xi_n)c(\xi')p_0c(\xi') + i\xi_nc(dx_n)p_0c(dx_n) \\
&\quad + (2 + i\xi_n)c(\xi')c(dx_n)\partial_{x_n}c(\xi') + ic(dx_n)p_0c(\xi') + ic(\xi')p_0c(dx_n) - i\partial_{x_n}c(\xi')] \\
&= \frac{1}{4(\xi_n - i)^2} \left[\frac{5}{2}h'(0)c(dx_n) - \frac{5i}{2}h'(0)c(\xi') - (2 + i\xi_n)c(\xi')c(dx_n)\partial_{\xi_n}c(\xi') + i\partial_{\xi_n}c(\xi') \right]; \tag{3.41}
\end{aligned}$$

$$B_2 = \frac{h'(0)}{2} \left[\frac{c(dx_n)}{4i(\xi_n - i)} + \frac{c(dx_n) - ic(\xi')}{8(\xi_n - i)^2} + \frac{3\xi_n - 7i}{8(\xi_n - i)^3} [ic(\xi') - c(dx_n)] \right]. \tag{3.42}$$

By (3.14), we obtain

$$\partial_{\xi_n} \sigma_{-3}(D^{-3}) = \frac{-4i\xi_nc(\xi') + (i - 3i\xi_n^2)c(dx_n)}{(1 + \xi_n^2)^3}. \tag{3.43}$$

By (3.27), (3.42) and (3.43), we have

$$\begin{aligned}
&\text{tr}[B_2 \times \partial_{\xi_n} \sigma_{-3}(D^{-3})(x_0)]|_{|\xi'|=1} \\
&= \text{tr} \left\{ \frac{h'(0)}{2} \left[\frac{c(dx_n)}{4i(\xi_n - i)} + \frac{c(dx_n) - ic(\xi')}{8(\xi_n - i)^2} + \frac{3\xi_n - 7i}{8(\xi_n - i)^3} [ic(\xi') - c(dx_n)] \right] \right. \\
&\quad \left. \times \frac{-4i\xi_nc(\xi') + (i - 3i\xi_n^2)c(dx_n)}{(1 + \xi_n^2)^3} \right\} \\
&= h'(0) \frac{4i - 11\xi_n - 6i\xi_n^2 + 3\xi_n^3}{(\xi_n - i)^5(\xi_n + i)^3}. \tag{3.44}
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
&\text{tr}[B_1 \times \partial_{\xi_n} \sigma_{-3}(D^{-3})(x_0)]|_{|\xi'|=1} \\
&= \text{tr} \left\{ \frac{1}{4(\xi_n - i)^2} \left[\frac{5}{2}h'(0)c(dx_n) - \frac{5i}{2}h'(0)c(\xi') - (2 + i\xi_n)c(\xi')c(dx_n)\partial_{\xi_n}c(\xi') + i\partial_{\xi_n}c(\xi') \right] \right. \\
&\quad \left. \times \frac{-4i\xi_nc(\xi') + (i - 3i\xi_n^2)c(dx_n)}{(1 + \xi_n^2)^3} \right\} \\
&= h'(0) \frac{3 + 12i\xi_n + 3\xi_n^2}{(\xi_n - i)^4(\xi_n + i)^3}. \tag{3.45}
\end{aligned}$$

Combining (3.44) and (3.45), we obtain

$$\begin{aligned}
\text{case c)} &= -ih'(0) \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{-7i + 26\xi_n + 15i\xi_n^2}{(\xi_n - i)^5(\xi_n + i)^3} d\xi_n \sigma(\xi') dx' \\
&= -ih'(0) \frac{2\pi i}{4!} \left[\frac{-7i + 26\xi_n + 15i\xi_n^2}{(\xi_n + i)^3} \right]^{(5)} \Big|_{\xi_n=i} \Omega_4 dx' \\
&= \frac{55}{16} \pi h'(0) \Omega_4 dx'. \tag{3.46}
\end{aligned}$$

Since Φ is the sum of the cases a), b) and c), so

$$\Phi = \left(\frac{15}{16} - \frac{35i}{16}\right)\pi h'(0)\Omega_4 dx'. \quad (3.47)$$

Now recall the Einstein-Hilbert action for manifolds with boundary [12][13][14],

$$I_{\text{Gr}} = \frac{1}{16\pi} \int_M \text{sdvol}_M + 2 \int_{\partial M} K \text{dvol}_{\partial M} := I_{\text{Gr},i} + I_{\text{Gr},b}, \quad (3.48)$$

where

$$K = \sum_{1 \leq i,j \leq n-1} K_{i,j} g_{\partial M}^{i,j}; \quad K_{i,j} = -\Gamma_{i,j}^n, \quad (3.49)$$

and $K_{i,j}$ is the second fundamental form, or extrinsic curvature. Taking the metric in Section 2, then by Lemma A.2 [12], for $n = 6$, then

$$K(x_0) = -\frac{5}{2}h'(0); \quad I_{\text{Gr},b} = -5h'(0)\text{Vol}_{\partial M}. \quad (3.50)$$

Then we obtain

Theorem 3.4. *Let M be a 6-dimensional compact spin manifold with the boundary ∂M and the metric g^M as above and D be the Dirac operator on $\widehat{M}$, then*

$$\text{Vol}_6^{(1,3)} = \widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{-3}] = -\frac{5\Omega_5}{3} \int_M \text{sdvol}_M + \left(\frac{7i}{8} - \frac{3}{8}\right)\pi\Omega_4 \int_{\partial M} K d\text{Vol}_{\partial M}. \quad (3.51)$$

Remark 3.5. *In [12] [13], Wang computed $\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{-1}]$ and $\widetilde{\text{Wres}}[\pi^+ D^{-2} \circ \pi^+ D^{-2}]$. In that cases, the boundary terms vanished, where the two operators are symmetric. Theorem 3.4 states the boundary terms is non-zero when we compute $\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{-3}]$. The reason is that D^{-1} and D^{-3} are not symmetric.*

Let

$$\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{-3}] = \widetilde{\text{Wres}}_i[\pi^+ D^{-1} \circ \pi^+ D^{-3}] + \widetilde{\text{Wres}}_b[\pi^+ D^{-1} \circ \pi^+ D^{-3}], \quad (3.52)$$

where

$$\widetilde{\text{Wres}}_i[\pi^+ D^{-1} \circ \pi^+ D^{-3}] = \int_M \int_{|\xi|=1} \text{trace}_{S(TM)}[\sigma_{-6}(D^{-1-3})]\sigma(\xi)dx \quad (3.53)$$

and

$$\begin{aligned} & \widetilde{\text{Wres}}_b[\pi^+ D^{-1} \circ \pi^+ D^{-3}] \\ = & \int_{\partial M} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{j,k=0}^{\infty} \sum \frac{(-i)^{|\alpha|+j+k+1}}{\alpha!(j+k+1)!} \times \text{trace}_{S(TM)}[\partial_{x_n}^j \partial_{\xi'}^\alpha \partial_{\xi_n}^k \sigma_r^+(D^{-1})(x', 0, \xi', \xi_n) \\ & \times \partial_x^\alpha \partial_{\xi_n}^{j+1} \partial_{x_n}^k \sigma_l(D^{-3})(x', 0, \xi', \xi_n)] d\xi_n \sigma(\xi') dx' \end{aligned} \quad (3.54)$$

denote the interior term and boundary term of $\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{-3}]$.

Combining (3.48), (3.51) and (3.52), we obtain

Corollary 3.6. *Let M be a 6-dimensional compact spin manifold with the boundary ∂M and the metric g^M as above and D be the Dirac operator on $\widehat{M}$, then*

$$\begin{aligned} I_{\text{Gr},i} &= \frac{-3}{80\pi\Omega_5} \widetilde{\text{Wres}}_i[\pi^+ D^{-1} \circ \pi^+ D^{-3}]; \\ I_{\text{Gr},b} &= \frac{16}{(7i-3)\pi\Omega_4} \widetilde{\text{Wres}}_b[\pi^+ D^{-1} \circ \pi^+ D^{-3}]. \end{aligned} \quad (3.55)$$

Nextly, for 5-dimensional spin manifolds with boundary, we compute $\text{Vol}_5^{(1,3)}$. By Proposition 2.2 (2) in [13], we have

$$\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{-3}] = \int_{\partial M} \Phi. \quad (3.56)$$

By (2.4), when $n = 5$, we have $r - k - |\alpha| + l - j - 1 = -5$, $r \leq -1, l \leq -3$, so we get $r = -1, l = -3, k = |\alpha| = j = 0$, then

$$\Phi = \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}_{S(TM)}[\sigma_{-1}^+(D^{-1})(x', 0, \xi', \xi_n) \times \partial_{\xi_n} \sigma_{-3}(D^{-3})(x', 0, \xi', \xi_n)] d\xi_n \sigma(\xi') dx'. \quad (3.57)$$

By (2.2.44) in [12], we have

$$\pi_{\xi_n}^+ \sigma_{-1}(D^{-1})(x_0)|_{|\xi'|=1} = \frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)}. \quad (3.58)$$

From (3.14) we obtain

$$\partial_{\xi_n} \sigma_{-3}(D^{-3}) = \frac{-4i\xi_n c(\xi') + (i - 3i\xi_n^2)c(dx_n)}{(1 + \xi_n^2)^3}. \quad (3.59)$$

Since $n = 5$, $\text{tr}(id) = \dim(S(TM)) = 4$. By the relation of the Clifford action and $\text{tr}AB = \text{tr}BA$, then we have the equalities:

$$\text{tr}[c(\xi')c(dx_n)] = 0; \quad \text{tr}[c(dx_n)^2] = -4; \quad \text{tr}[c(\xi')^2](x_0)|_{|\xi'|=1} = -4. \quad (3.60)$$

Hence from (3.58), (3.59), and (3.60), we have

$$\begin{aligned} & \text{tr} \left[\left(\frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)} \right) \times \left(\frac{-4i\xi_n c(\xi') + (i - 3i\xi_n^2)c(dx_n)}{(1 + \xi_n^2)^3} \right) \right] \\ &= \frac{2i - 6\xi_n}{(\xi_n - i)^3(\xi_n + i)^3}. \end{aligned} \quad (3.61)$$

Then

$$\begin{aligned} \Phi &= \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{2i - 6\xi_n}{(\xi_n - i)^3(\xi_n + i)^3} d\xi_n \sigma(\xi') dx' \\ &= \frac{2\pi i}{2!} \left[\frac{2i - 6\xi_n}{(\xi_n + i)^3} \right]^{(2)} \Big|_{\xi_n=i} \Omega_3 dx' \\ &= \frac{3\pi i}{4} \Omega_3 dx'. \end{aligned} \quad (3.62)$$

By $\text{Vol}_5^{(1,3)} = \frac{3\pi i}{4} \Omega_3 \text{Vol}_{\partial M}$ and $I_{\text{Gr,b}} = -4h'(0) \text{Vol}_{\partial M}$, we have

Theorem 3.7. *Let M be a 5-dimensional compact spin manifold with the boundary ∂M and the metric g^M as in Section 2 and D be the Dirac operator on $\widehat{M}$, then*

$$\text{Vol}_5^{(1,3)} = \widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{-3}] = \frac{3\pi i}{4} \Omega_3 \text{Vol}_{\partial M}, \quad (3.63)$$

$$I_{\text{Gr,b}} = \frac{16ih'(0)}{3\pi\Omega_3} \widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{-3}], \quad (3.64)$$

where $\text{Vol}_{\partial M}$ denotes the canonical volume of ∂M .

4. A Kastler-Kalau-Walze type theorem for perturbations of Dirac operators

In [15], Connes and Moscovici defined a twisted spectral triple and considered the perturbations of Dirac operator $e^h D e^h$. In this section, for perturbations of Dirac operators, we compute the lower dimensional volume $\text{Vol}_6^{(2,2)}$ for 6-dimensional spin manifolds with boundary and get a Kastler-Kalau-Walze type theorem in this case.

4.1. $\widetilde{\text{Wres}}[\pi^+(fD^{-2}) \circ \pi^+D^{-2}]$ for the Dirac operators

Let M be a 6-dimensional compact spin manifold with the boundary ∂M and the metric g^M as Section 2 and D be the Dirac operator on $\widehat{M}$, we will compute $\widetilde{\text{Wres}}[\pi^+(fD^{-2}) \circ \pi^+D^{-2}]$ for a smooth function $f(x)$. By the Kastler-Kalau-Walze type theorem in [6], we get the lower dimensional volume for 6-dimensional spin manifolds without boundary. An application of Theorem 1 in [13] shows that

Lemma 4.1. *Let M be a 6-dimensional compact spin manifold without boundary, then*

$$\text{Wres}[fD^{-4}] = -\frac{5\Omega_5}{3} \int_M f \text{sdvol}_M. \quad (4.1)$$

Therefore, we only need to compute $\int_{\partial M} \Phi$. By Lemma 1 in [13], we have

Lemma 4.2. *For 6-dimensional compact spin manifold with the boundary ∂M and the metric g^M as above, then*

$$\sigma_{-2}(D^{-2}) = |\xi|^{-2}; \quad (4.2)$$

$$\sigma_{-2}(fD^{-2}) = f|\xi|^{-2}; \quad (4.3)$$

$$\sigma_{-3}(D^{-2}) = -\sqrt{-1}|\xi|^{-4}\xi_k(\Gamma^k - 2\delta^k) - \sqrt{-1}|\xi|^{-6}2\xi^j\xi_\alpha\xi_\beta\partial_j g^{\alpha\beta}. \quad (4.4)$$

Since Φ is a global form on ∂M , so for any fixed point $x_0 \in \partial M$, we can choose the normal coordinates U of x_0 in ∂M (not in M) and compute $\Phi(x_0)$ in the coordinates $\widetilde{U} = U \times [0, 1) \subset M$ and the metric $\frac{1}{h(x_n)}g^{\partial M} + dx_n^2$. For details, see Section 2.2.2[12]. Now we can compute Φ (see formula (2.4) for the definition of Φ), since the sum is taken over $-r-l+k+j+|\alpha|=5$, $r, l \leq -2$, then we have the following five cases:

case a) I) $r = -2, l = -2, k = j = 0, |\alpha| = 1$

From (2.4) we have

$$\text{case a) I)} = -f \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{|\alpha|=1} \text{trace}[\partial_{\xi'}^\alpha \pi_{\xi_n}^+ \sigma_{-2}(D^{-2}) \times \partial_{\xi_n}^\alpha \sigma_{-2}(D^{-2})](x_0) d\xi_n \sigma(\xi') dx'. \quad (4.5)$$

By Lemma 3.2, for $i < n$, then

$$\partial_{x_i} \sigma_{-2}(D^{-2})(x_0) = \partial_{x_i}(|\xi|^{-2})(x_0) = -\frac{\partial_{x_i}(|\xi|^2)(x_0)}{|\xi|^4} = 0. \quad (4.6)$$

Then case a) I) vanishes.

case a) II) $r = -2, l = -2, k = |\alpha| = 0, j = 1$

From (2.4) we have

$$\begin{aligned} \text{case a) II)} &= -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_{-2}(fD^{-2}) \times \partial_{\xi_n}^2 \sigma_{-2}(D^{-2})](x_0) d\xi_n \sigma(\xi') dx' \\ &= -\frac{1}{2} f \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_{-2}(D^{-2}) \times \partial_{\xi_n}^2 \sigma_{-2}(D^{-2})](x_0) d\xi_n \sigma(\xi') dx' \\ &\quad - \frac{1}{2} \partial_{x_n}(f) \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\pi_{\xi_n}^+ \sigma_{-2}(D^{-2}) \times \partial_{\xi_n}^2 \sigma_{-2}(D^{-2})](x_0) d\xi_n \sigma(\xi') dx'. \end{aligned} \quad (4.7)$$

By **case a) II)** in [13], we have

$$-\frac{1}{2} f \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_{-2}(D^{-2}) \times \partial_{\xi_n}^2 \sigma_{-2}(D^{-2})](x_0) d\xi_n \sigma(\xi') dx' = -\frac{5}{8} \pi f h'(0) \Omega_4 dx', \quad (4.8)$$

where Ω_4 is the canonical volume of S^4 .

On the other hand, by (14) in [13], we have

$$\partial_{\xi_n}^2 (\sigma_{-2}(D^{-2}))(x_0) = \partial_{\xi_n}^2 (|\xi|^{-2})(x_0) = \frac{-2 + 6\xi_n^2}{(1 + \xi_n^2)^3}. \quad (4.9)$$

By (4.2) and the Cauchy integral formula, then

$$\begin{aligned} \pi_{\xi_n}^+ \sigma_{-2}(D^{-2})(x_0)|_{|\xi'|=1} &= \frac{1}{2\pi i} \lim_{u \rightarrow 0^-} \int_{\Gamma^+} \frac{1}{(\eta_n + i)(\xi_n + iu - \eta_n)} d\eta_n \\ &= \frac{-i}{2(\xi_n - i)}. \end{aligned} \quad (4.10)$$

Combining (4.9) and (4.10), we have

$$\text{tr} \left[\frac{-i}{2(\xi_n - i)} \times \frac{-2 + 6\xi_n^2}{(1 + \xi_n^2)^3} \right] = \frac{i - 3i\xi_n^2}{(\xi_n - i)^4(\xi_n + i)^3} \text{tr}[id] = \frac{8i - 24i\xi_n^2}{(\xi_n - i)^4(\xi_n + i)^3}, \quad (4.11)$$

where $n = 6$, $\text{tr}_{S(TM)}[id] = \dim(\wedge^*(3)) = 8$.

From (4.7) and (4.11), we have

$$\begin{aligned} & -\frac{1}{2} \partial_{x_n} f \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\pi_{\xi_n}^+ \sigma_{-2}(D^{-2}) \times \partial_{\xi_n}^2 \sigma_{-2}(D^{-2})](x_0) d\xi_n \sigma(\xi') dx' \\ &= -\frac{1}{2} \partial_{x_n} (f) \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{8i - 24i\xi_n^2}{(\xi_n - i)^4(\xi_n + i)^3} (x_0) d\xi_n \sigma(\xi') dx' \\ &= -\frac{1}{2} \partial_{x_n} (f) \frac{2\pi i}{3!} \left[\frac{8i - 24i\xi_n^2}{(\xi_n + i)^3} \right]^{(3)}|_{\xi_n=i} \Omega_3 dx' \\ &= 3\pi i \partial_{x_n} (f) \Omega_4 dx'. \end{aligned} \quad (4.12)$$

Hence in this case,

$$\text{case a) II) } = -\frac{5}{8} \pi f h'(0) \Omega_4 dx' + 3\pi i \partial_{x_n} (f) \Omega_4 dx'. \quad (4.13)$$

case a) III) $r = -2$, $l = -2$ $j = |\alpha| = 0$, $k = 1$

From (2.4) we have

$$\begin{aligned} \text{case a) III) } &= -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-2}(f D^{-2}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-2}(D^{-2})](x_0) d\xi_n \sigma(\xi') dx' \\ &= -\frac{1}{2} f \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-2}(D^{-2}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-2}(D^{-2})](x_0) d\xi_n \sigma(\xi') dx'. \end{aligned} \quad (4.14)$$

By **case a) III)** in [13], we have

$$-\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-2}(D^{-2}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-2}(D^{-2})](x_0) d\xi_n \sigma(\xi') dx' = \frac{5}{8} \pi h'(0) \Omega_4 dx'. \quad (4.15)$$

Combining (4.14) and (4.15), we obtain

$$\text{case a) III) } = \frac{5}{8} \pi f h'(0) \Omega_4 dx'. \quad (4.16)$$

Since **case b)**, **case c)** has the same expression with the case of **case b)**, **case c)** in [13] multiplied by a function f , so we can use the same way to compute the two terms. An application of (21) and (24) in [13] shows that the sum of **case b)** and **case c)** is zero. Hence we conclude that, the sum of **case a)**, **case b)** and **case c)** is

$$\Phi = 3\pi i \partial_{x_n} (f) \Omega_4 dx'. \quad (4.17)$$

In summary, we have proved

Theorem 4.3. *Let M be a 6-dimensional compact spin manifold with the boundary ∂M and the metric g^M as above and D be the Dirac operator on $\widehat{M}$, then*

$$\widetilde{\text{Wres}}[\pi^+(fD^{-2}) \circ \pi^+D^{-2}] = -\frac{5\Omega_5}{3} \int_M f \text{sdvol}_M + 3\pi i \Omega_4 \int_{\partial M} \partial_{x_n}(f)|_{x_n=0} \text{dvol}_{\partial M}. \quad (4.18)$$

4.2. $\widetilde{\text{Wres}}[\pi^+(D^2 + f)^{-1} \circ \pi^+D^{-2}]$ for the Dirac operators

In this subsection, let M be a 6-dimensional compact spin manifold with the boundary ∂M and the metric g^M as Section 2 and D be the Dirac operator on $\widehat{M}$, we will compute $\widetilde{\text{Wres}}[(D^2 + f)^{-1}D^{-2}]$ for smooth function $f(x)$.

Firstly, we compute the symbol $\sigma((D^2 + f)^{-1})$ of $(D^2 + f)^{-1}$. By the definition of the Dirac operator D in section 3 and Lemma 1 in [13], we have

$$\sigma_{-2}((D^2 + f)^{-1}) = \sigma_{-2}(D^{-2}); \quad \sigma_{-3}((D^2 + f)^{-1}) = \sigma_{-3}(D^{-2}). \quad (4.19)$$

Let $\tilde{D} = D^4 + fD^2$ be the fourth order operators. Write

$$D_x^\alpha = (-\sqrt{-1})^{|\alpha|} \partial_x^\alpha; \quad \sigma(\tilde{D}) = \tilde{p}_4 + \tilde{p}_3 + \tilde{p}_2 + \tilde{p}_1 + \tilde{p}_0; \quad \sigma((\tilde{D})^{-1}) = \sum_{j=4}^{\infty} \tilde{q}_{-j}. \quad (4.20)$$

Since the smooth function $f(x) \in \tilde{p}_0$, then we have

$$\sigma_4(\tilde{D}^4) = \sigma_4(D^4); \quad \sigma_3(\tilde{D}^4) = \sigma_3(D^4); \quad \sigma_2(\tilde{D}^4) = \sigma_2(D^4) + f|\xi|^2. \quad (4.21)$$

By the composition formula of pseudodifferential operators, we have

$$\begin{aligned} 1 = \sigma(\tilde{D} \circ \tilde{D}^{-1}) &= \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} [\sigma(\tilde{D})] D_x^{\alpha} [\sigma((\tilde{D})^{-1})] \\ &= (\tilde{p}_4 + \tilde{p}_3 + \tilde{p}_2 + \tilde{p}_1 + \tilde{p}_0)(\tilde{q}_{-4} + \tilde{q}_{-5} + \tilde{q}_{-6} + \cdots) \\ &\quad + \sum_j (\partial_{\xi_j} \tilde{p}_4 + \partial_{\xi_j} \tilde{p}_3 + \partial_{\xi_j} \tilde{p}_2 + \partial_{\xi_j} \tilde{p}_1 + \partial_{\xi_j} \tilde{p}_0) \\ &\quad \times (D_{x_j} \tilde{q}_{-4} + D_{x_j} \tilde{q}_{-5} + D_{x_j} \tilde{q}_{-6} + \cdots) \\ &\quad + \sum_{i,j} (\partial_{\xi_i} \partial_{\xi_j} (\tilde{p}_4 + \tilde{p}_3 + \tilde{p}_2 + \tilde{p}_1 + \tilde{p}_0)) \\ &\quad \times (D_{x_i} D_{x_j} \tilde{q}_{-4} + D_{x_i} D_{x_j} \tilde{q}_{-5} + D_{x_i} D_{x_j} \tilde{q}_{-6} + \cdots) \\ &= \tilde{p}_4 \tilde{q}_{-4} + (\tilde{p}_4 \tilde{q}_{-5} + \tilde{p}_3 \tilde{q}_{-4} + \sum_j \partial_{\xi_j} \tilde{p}_4 D_{x_j} \tilde{q}_{-4}) + \cdots. \end{aligned} \quad (4.22)$$

Then we obtain

$$\tilde{q}_{-4} = \tilde{p}_4^{-1}; \quad (4.23)$$

$$\tilde{q}_{-5} = -\tilde{p}_4^{-1} [\tilde{p}_3 \tilde{q}_{-4} + \sum_j \partial_{\xi_j} \tilde{p}_4 D_{x_j} (\tilde{q}_{-4})]; \quad (4.24)$$

$$\begin{aligned} 0 &= \tilde{p}_4 \tilde{q}_{-6} + \tilde{p}_3 \tilde{q}_{-5} + \tilde{p}_2 \tilde{q}_{-4} + \sum_j \partial_{\xi_j} \tilde{p}_3 D_{x_j} \tilde{q}_{-4} + \sum_j \partial_{\xi_j} \tilde{p}_4 D_{x_j} \tilde{q}_{-5} \\ &\quad + \sum_{i,j} \partial_{\xi_i} \partial_{\xi_j} \tilde{p}_4 D_{x_i} D_{x_j} \tilde{q}_{-4}. \end{aligned} \quad (4.25)$$

Hence from (4.21)-(4.25) and the recursion formula, we have

Lemma 4.4. *For a 6-dimensional compact spin manifold with the boundary ∂M , then*

$$q_{-6} = -f|\xi|^2 + \sigma_{-6}(D^{-4}). \quad (4.26)$$

By Lemma 4.4 and [5], [6], we obtain

Theorem 4.5. *let M be a 6-dimensional compact spin manifold without boundary and $f_i (i = 1, 2, 3)$ be the smooth function, then*

$$\text{Wres}[(D^4 + f_1 D^2 + f_2 D + f_3)^{-1}] = -\Omega_5 \int_M (\frac{5}{3}s + 8f_1) \text{dvol}_M. \quad (4.27)$$

Remark 4.6. *When $f_i = 0$ ($i = 1, 2, 3$), we get the classical Kastler-Kalau-Walze type theorem.*

On the other hand, it is straightforward to see from (4.19) we can directly get the same results with Theorem 1 in [13]. Hence we conclude that $\int_{\partial M} \Phi = 0$. In summary, we have proved

Theorem 4.7. *let M be a 6-dimensional compact spin manifold with the boundary ∂M and the metric g^M as Section 2 and D be the Dirac operator on $\bar{M}$, then*

$$\widetilde{\text{Wres}}[\pi^+(D^2 + f)^{-1} \circ \pi^+ D^{-2}] = -\Omega_5 \int_M (\frac{5}{3}s + 8f) \text{dvol}_M. \quad (4.28)$$

Acknowledgements

This work was supported by Fok Ying Tong Education Foundation under Grant No. 121003 and NSFC. 11271062. The author also thank the referee for his (or her) careful reading and helpful comments.

References

References

- [1] V. W. Guillemin.: A new proof of Weyl's formula on the asymptotic distribution of eigenvalues. Adv. Math. 55, no. 2, 131-160, (1985).
- [2] M. Wodzicki.: local invariants of spectral asymmetry. Invent. Math. 75(1), 143-178, (1995).
- [3] A. Connes.: Quantized calculus and applications. XIth International Congress of Mathematical Physics(Paris,1994), Internat Press, Cambridge, MA, 15-36, (1995).
- [4] A. Connes.: The action functional in Noncommutative geometry. Comm. Math. Phys. 117, 673-683, (1998).
- [5] D. Kastler.: The Dirac Operator and Gravitation. Comm. Math. Phys. 166, 633-643, (1995).
- [6] W. Kalau and M. Walze.: Gravity, Noncommutative geometry and the Wodzicki residue. J. Geom. Physics. 16, 327-344,(1995).
- [7] T. Ackermann.: A note on the Wodzicki residue. J. Geom.Phys. 20, 404-406, (1996).
- [8] R. Ponge.: Noncommutative geometry and lower dimensional volumes in Riemannian geometry. Letters in Mathematical Physics. 83, no.1, 19-32, (2008).
- [9] B. V. Fedosov, F. Golse, E. Leichtnam, E. Schrohe.: The noncommutative residue for manifolds with boundary. J. Funct. Anal. 142, 1-31, (1996).
- [10] Y. Wang.: Differential forms and the Wodzicki residue for Manifolds with Boundary. J. Geom. Physics. 56, 731-753, (2006).
- [11] Y. Wang.: Differential forms the Noncommutative Residue for Manifolds with Boundary in the non-product Case, Letters in Mathematical Physics. 77, 41-51, (2006).
- [12] Y. Wang.: Gravity and the Noncommutative Residue for Manifolds with Boundary. Letters in Mathematical Physics. 80, 37-56, (2007).
- [13] Y. Wang.: Lower-Dimensional Volumes and Kastler-kalau-Walze Type Theorem for Manifolds with Boundary . Commun. Theor. Phys. Vol 54, 38-42, (2010).
- [14] Y. Yu.: The Index Theorem and The Heat Equation Method, Nankai Tracts in Mathematics-Vol.2, World Scientific Publishing, (2001).
- [15] A. Connes, H. Moscovici.: Type III and spectral. Traces in Geometry, Number Theory and Quantum Fields, Aspects of Mathematics E38, Vieweg Verlag, 57-71, (2008).